

PROSPECTIVE RANDOMISED CONTROL STUDY OF POST OP EPIDURAL ANALGESIA WITH BUPIVACAINE AND FENTANYL VS. BUPIVACAINE AND CLONIDINERachana N. D¹, Saraswathi N², Srinivas Babu A³, Anup N⁴**HOW TO CITE THIS ARTICLE:**

Rachana N. D, Saraswathi N, Srinivas Babu A. Anup N. "Prospective Randomised Control Study of Post Op Epidural Analgesia with Bupivacaine and Fentanyl Vs. Bupivacaine and Clonidine". Journal of Evolution of Medical and Dental Sciences 2014; Vol. 3, Issue 31, July 31; Page: 8704-8711, DOI: 10.14260/jemds/2014/3110

ABSTRACT: The aim of the study was to compare the relative potencies and clinical characteristics of epidural Clonidine and Fentanyl with Bupivacaine in lower limb and lower abdominal surgeries using patient-controlled analgesia. In a randomised double-blinded study, 60 ASA I or II patients requiring epidural analgesia for post-operative pain relief were allocated to receive either 0.125% Bupivacaine with Clonidine 1µg/kg or 0.125% Bupivacaine with Fentanyl 1µg/kg via a sterile syringe by trained anaesthesiologists. Analgesia was established with 10-15 ml bolus of study solution. There were significant differences in onset time, duration and quality of analgesia, local anaesthetic consumption, between the two groups. We conclude that 0.125% Bupivacaine with Clonidine 1µg/kg group of patients clinically had better quality of analgesia and for a longer duration in comparison with patients receiving 0.125% Bupivacaine with Fentanyl 1µg/kg.

KEYWORDS: Patient-controlled analgesia, epidural anaesthesia, post-operative pain.

INTRODUCTION: Pain after surgery is inevitable, yet it is often underestimated and undertreated. Besides the distress, pain is also associated with psychosocial effects. The effective relief of pain is of paramount importance to anyone treating patients undergoing surgery. This has a smoother postoperative course with early discharge and also reduces the chronic pain syndromes.¹

In our setup, epidural Bupivacaine is the most widely used local anaesthetic agent for post-operative epidural analgesia. Fentanyl is frequently used in combination with Bupivacaine, as its use permits effective analgesia using concentrations of Bupivacaine that would otherwise be sub-therapeutic. In the last decade, so many other drugs have been used as an adjuvant to local anaesthetic agents, to improve the efficacy and potency of local anaesthetic agents and to provide good analgesia.

Epidural technique has been found to provide better pain relief than systemic opioids and also decreased incidence of postoperative complications. Lumbar epidural catheters can be kept in place for prolonged periods. Epidural administration of various analgesics gained increasing popularity following the discovery of opioids receptors in the spinal cord capable of producing potent analgesia as reported by Taksh and Rudy in 1976.⁽²⁾ The epidural analgesia does inhibit the stress response. This effect seems to be greatest when epidural anaesthesia is continued in the post-operative period as epidural analgesia.

The current trend in postoperative pain management is multimodal analgesia. Experimentally, Clonidine, local anesthetics and opioids interact synergistically and clinically, are at least additive.

The analgesic neuraxial effect of the α₂-adrenergic agonist Clonidine is well described⁽³⁾ Epidural Clonidine improved analgesia when combined with epidural Bupivacaine- Fentanyl in

ORIGINAL ARTICLE

patients after lower abdominal surgery⁽⁴⁾ The α 2-adrenergic agonist Clonidine produces analgesia by different mechanism that mimics the effect of endogenously released Nor-Epinephrine to stimulate postsynaptic α 2 receptors in the spinal cord.

Widespread application of opioids in spinal analgesia has been limited by side effects such as life threatening and unpredictable respiratory depression. Hence α 2 adrenergic agonist, Clonidine is being used as an alternative drug for postoperative analgesia which is free of some opioids related but not all side effects.

METHODS: Following Local Research Ethics Committee approval and written informed consent, 60 ASA I or II patients aged 18-75 yrs who were undergoing lower abdominal and lower limb surgeries were included in our study.

Exclusion Criteria Included: contraindication to epidural anesthesia; sensitivity or allergy to amide local anesthetics; coexisting disease which could affect the reliability of clinical assessments; known or suspected drug abuse; and pregnancy, proven or possible. Preoperatively, each patient provided a medical history and underwent routine physical examination. The investigator or research assistant instructed patients in the use of epidural analgesia and the visual analog scale (VAS) pain score.

Oral or parenteral benzodiazepines could be administered preoperatively at the discretion of the investigator. After surgery patients were randomised to one of the study groups.

Prior to surgery an epidural catheter was inserted in a lower thoracic or upper lumbar intervertebral space and threaded cephalad 4 to 5 cm. Patients received lidocaine 2% 3 mL with added epinephrine 5 μ g/mL as a catheter test dose, followed by study drugs 10-20 mL over 5mins.

In the post-anesthesia recovery room, when fully awake, patient was asked to score the pain on the basis of VAS. The post-operative epidural bolus was given when the patients complains of moderate pain. In group-A patients, epidural bolus of 0.125% Bupivacaine (10-15ml) + Clonidine (1 μ g/kg) will be delivered whenever patient complains of pain at a minimum interval of 4 hours. In group-B patients, a combination of 0.125%Bupivacaine (10-15ml) + Fentanyl (1 μ g/kg) will be given at similar interval. Other analgesics given during the study period were noted.

In the post-operative period the following parameters were studied: Vital parameters such as the heart rate, blood pressure, respiratory rate, sedation score and visual analogue score were recorded. Duration of analgesia: This was calculated from the time when the first dose of analgesia was given, postoperatively and followed up till the patient complained of pain. Patients were asked to point out the intensity of their pain on the visual analogue scale.

Time at which patients complained of pain more than 4 on the scale was noted. That point was taken as the end of fair analgesia and at that point, top up doses of same drug with same dosage given. And the number of additional doses (top up doses) required in first 24 hrs were noted in both the groups. Side effects: Like nausea, vomiting, hypotension, respiratory depression and pruritus allergic reaction were looked for.

Motor blockade assessed according to a four-point modified Bromage scale (0 = no motor block, 1 = inability to raise extended leg, 2 = inability to flex knee, and 3 =inability to flex ankle joint). Sedation was assessed as Score: 0= alert; 1=drowsy;2=sleeps, easy to arouse verbally, does not fall asleep during or immediately after conversation, can stand up; 3= sleeps, opens the eyes to verbal

command, falls asleep during or immediately after conversation, cannot stand up; 4= does not open the eyes to verbal command.

Patients were instructed to inform when pain of any intensity occurred and pain scoring was done according to Visual Analogue Scale. Epidural bolus as mentioned above will be administered at VAS of 4 or more. Each time the bolus was given parameters mentioned above were recorded every 15 minutes initially for first hour then afterwards hourly till the next bolus was given for 24 hours postoperatively.

Descriptive statistical analysis has been carried out in the present study. Results on continuous measurements are presented on Mean \pm SD (Min-Max) and results on categorical measurements are presented in Number (%). Significance is assessed at 5 % level of significance. The following assumptions on data are made, Assumptions:

1. Dependent variables should be normally distributed,
2. Samples drawn from the population should be random, and Cases of the samples should be independent. Student's T test was used to find the significance of the study parameters on the continuous scale, Inter group analysis on metric parameters and chi square test to find the significance of parameters on categorical scale between two or more groups.

RESULTS: In our study 60 patients of ASA I and II were included in the study and were allocated to the study groups randomly. In our study patients in both the groups belonged to age group between 31-40 years and there was no significant difference in the sex distribution and anthropometry. In our study we found that SpO₂ was significantly stable in group A (99.45—99.20) patients compared to group B (98.78 – 99.08) patients. (Graph 1) Group A patients had statistically significant bradycardia with P<0.001 (graph 2). Group A patients had significant hypotension P<0.001. (Graph 3A and 3B).

Analysis of severity of pain was assessed based on VAS scores and quality of analgesia. (graph 4) Group A patients had better quality of analgesia with faster onset and analgesia lasting for 6-8 hours with less number of bolus doses, Group A patients needed less number of bolus doses (4 Doses) than the group B patients (6 doses). The Need for rescue analgesics was less in group-A patients (93.3% needed no rescue analgesics) P=0.254 which again shows that quality of analgesia was better in group A patients. (TABLE 1) The adverse effects were also noted in the patients during our study.

We found that Group A patients had hypotension (12%), sedation (20%), and transient motor blockade (12%), while nausea (33%) and pruritus (16.7%) were most common with group B patients. (TABLE 2).

DISCUSSION: In the postoperative period when the effect of the anaesthetic disappears, the tissue injury persists and pain producing substances which are liberated during the operation greatly reduce the normally high threshold of the nociceptors, so that innocuous stimulation produces pain.

Moreover the cut ends of axons further contribute to nociception. Surgical trauma and subsequent postoperative pain result in a broad range of endocrinological, immunological and inflammatory responses, including increased release of catabolic hormones and inhibited secretion of anabolic mediators. To minimize or overcome these adverse effects, the postoperative pain should be optimally treated.

ORIGINAL ARTICLE

In this study, group-A patients received a combination of 0.125% Bupivacaine (10-15ml) +Clonidine (1µg/kg) and group-B patients received epidural bolus of 0.125% Bupivacaine (10-15ml) +Fentanyl (1µg/kg). Pain was measured using visual analogue scale.

This study clearly demonstrates that combination therapy with epidural Clonidine and Bupivacaine gives superior pain relief than combination of Fentanyl and Bupivacaine. Although several studies have reported the beneficial effect of epidural Clonidine for intra and post-operative pain management.^(5,6,7) This study is unique in comparing bolus dosing combination of Clonidine and Bupivacaine with Fentanyl and Bupivacaine.

The onset of analgesia was faster and duration of analgesia was prolonged significantly in Clonidine group as evidenced by VAS score measurement in comparison with Fentanyl group. Analgesia lasted for 6 hours in group-A where as in group-B it lasted for 4 hours and number of boluses required in first 24 hrs was less in Clonidine group.^{(5-6),(8-12),(13-16)}

In our study it was found that hypotension and bradycardia was seen more commonly with Clonidine group of patients at different time intervals but both hypotension and bradycardia did not require any medical intervention which further proves that Clonidine though causes hypotension and bradycardia there was no hemodynamic instability. In the Fentanyl group, there were no significant hemodynamic changes noted as compared to Clonidine group.^{(9),(4),(7),(17-19),(25)}

In the current study, drop in the oxygen saturation was very negligible in Clonidine as compared to Fentanyl group at all intervals. Even in Fentanyl group, although there was statistically significant drop in the Spo₂, it was maintained at or above 98 % at all intervals. So, there was no significant respiratory depression in both the groups.^(20,21)

In our study it was found that rescue analgesics were more needed in Fentanyl (20%) group of patients than compared to Clonidine(6.7%) group of Patients.^(5,9)

In our study, Clonidine was found to cause more sedation and alter the heart rate and blood pressure. Excess sedation was noted in 5 out of 30 patients in our study.^{(22-24),(16), (19).}

To conclude, In the management of postoperative pain relief for various surgical procedures both Clonidine and Fentanyl can be used safely and effectively as adjuvant to Bupivacaine. It can be concluded from the above study that epidural Clonidine has a better analgesic action and longer duration of analgesia in comparison to Fentanyl and also Clonidine group of patients required less bolus doses.

However the side effects such as hypotension, sedation and motor blockade were common in Clonidine whereas nausea vomiting, Pruritus were common in Fentanyl group. The overall side effects were less in Clonidine group as compared to Fentanyl group.

REFERENCES:

1. Dr. Ed Charlton. The management of postoperative pain. "Update in anaesthesia".1997; Issue 7.
2. Taksh TL, Rudy TA et al. An inability to antagonize with Naloxone the elevated nociceptive thresholds resulting from electrical stimulation of the mesencephalic central gray. Life Sci. 1976; 18:1193-8.
3. Eisenach, James C et al. α 2-Adrenergic Agonists for Regional Anesthesia: A Clinical Review of Clonidine (1984 - 1995). Anesthesiology: September 1996 - Volume 85 - Issue 3 - pp 655-674.


ORIGINAL ARTICLE

4. M Peach, T Pavy et al. Patient-controlled epidural analgesia in labor: The addition of Clonidine to Bupivacaine-Fentanyl. *Regional Anesthesia and Pain Medicine*;2000, Volume 25, Issue 1, 25: 31–40.
5. Kizilarслан S, Kuvaki B et al. Epidural fentanyl-bupivacaine compared with clonidine-bupivacaine for analgesia in labour. *European Journal of Anaesthesiology*: November 2000 - Volume 17 - Issue 11 - pp 692-69.
6. Lt. Col. Upadhyay K. K. Brig. Prabhakar T et al. Study of the efficacy and safety of clonidine as an adjunct to bupivacaine for caudal analgesia in children. *Indian J. anaesth*, 2005; 49. 3; 199-201.
7. Dobrydnjov I, Axelsson K, Gupta A, et al. Improved analgesia with clonidine when added to local anesthetic during combined spinal-epidural anesthesia for hip arthroplasty: a double blind, randomized and placebo controlled study. *Acta Anaesthesiol Scan* 2005; 49: 538–45.
8. Rostaing S, Bonnet F, Levron JC et al. Effect of epidural clonidine on analgesia and pharmacokinetics of epidural fentanyl in postoperative patients. *Anesthesiology*. 1991 Sep; 75 (3): 420-5.
9. Shobhana Gupta, Dipak Ravaletal. Addition of epidural Clonidine enhances postoperative analgesia: A double-blind study in total knee replacement surgeries. *Anesth essays res* 2010.4:2. 70-4.
10. Rawal N, Arner S, et al. Present state of extradural and intrathecal opioid analgesia in Sweden. *British Journal of Anaesthesia* 1987; 59: 791-799.
11. Kalia PK, Madan R, et al. Clinical study on epidural clonidine for postoperative analgesia. *Indian Journal of Medical Research* 1986; 83:550
12. Yuan-Shiou Huang, Liu-Chi Lin et al. Epidural Clonidine for Postoperative Pain after Total Knee Arthroplasty: A Dose-Response Study. *Anesth Analg* 2007; 104: 1230 –5.
13. Welchew E. A., D. P. Breen et al. Patient-controlled on-demand epidural fentanyl A comparison of patient-controlled on-demand fentanyl delivered epidurally or intravenously. *Anaesthesia*.1991. Volume 46, Issue 6, pages 438–441.
14. Carabine UA, Milligan KR, Moore J. Extradural clonidine and bupivacaine for postoperative analgesia. *Br J Anaesth* 1992; 68: 132–5.
15. M J Paech, T J Pavy, C E Orlikowski, W Lim, and S F Evans. Postoperative epidural infusion: a randomized, double-blind, dose-finding trial of clonidine in combination with bupivacaine and Fentanyl. *AnesthAnalg* June 1997 84:1323-1328.
16. Sveticic et al. Combinations of bupivacaine, fentanyl, and clonidine for lumbar epidural postoperative analgesia: a novel optimization procedure. *Anesthesiology* 2004; 101:1381–93.
17. Michael G. Rockemann, MD, Wulf Seeling et al. Analgesic and Hemodynamic Effects of Epidural Clonidine, Clonidine/Morphine, and Morphine After Pancreatic Surgery-A Double-Blind Study. *AnesthAnalg*1995; 80: 869-74.
18. Eisenach J, Detweiler D, et al. Hemodynamic and analgesic actions of epidurally administered clonidine. *Anesthesiology* 1993; 78: 277-87.
19. Ahmed Aly Fawaz. Extradural Clonidine versus Fentanyl for Breakthrough Pain in Advanced Vaginal Delivery in Ain Shams *Journal of Anesthesiology*. Jan 2012; Vol. 5-1:23-31
20. Bailey PL et al Respiratory effects of clonidine alone and combined with morphine, in humans. *Anesthesiology* 1991; 74: 43–8.

ORIGINAL ARTICLE


21. Negre I, Gueneron JP, Ecoffey C, Penon C, Gross JB, Levron JC, Samii K. Ventilatory response to carbon dioxide after intramuscular and epidural fentanyl. *Anesth Analg* 1987; 66: 707–10.
22. Armand S et al. Meta-analysis of the efficacy of extradural Clonidine to relieve postoperative pain *Br J Anaesth* 1998; 81: 126–34.
23. Cooper DW et al Patient-controlled extradural analgesia to compare bupivacaine, fentanyl and bupivacaine with fentanyl in the treatment of postoperative pain. *Br J Anaesth* 1993; 70: 503–7
24. Cucchiaro G et al Side effects of postoperative epidural analgesia in children: a randomized study comparing morphine and clonidine. *Paediatr Anaesth* 2003; 13: 318–23.
25. George KA, Chisakuta AM, Gamble JA, Browne GA. Thoracic epidural analgesia for postoperative pain relief following abdominal aortic surgery. *Anaesthesia* 1992; 47: 388-9.

Graph 1: Comparison of SPO₂ between two groups of patients studied which shows that group A (Clonidine) patients had no significant drop in saturations when compared to group B (Fentanyl) patients.



Graph 1


Graph 2: Comparison of Heart rate (bpm) between two groups of patients studied showed that group A patients had bradycardia in comparison to group B patients at various time intervals.



Graph 2


ORIGINAL ARTICLE

Graph 3A: Comparison of SBP (mm Hg) between two groups of patients studied.




Graph 3A

Graph 3B: Comparison of DBP (mm Hg) between two groups of patients studied showed that drop in blood pressures both SBP and DBP noticed with Group A patients.



Graph 3B

Graph 4: Comparison of Pain Score between two groups of patients studied that group A patients had better quality of analgesia assessed according to pain scores and analgesia lasted for slightly prolonged duration in comparison to group B patients.



Graph 4

Rescue analgesia	Group A (n=30)	Group B (n=30)
No	28(93.3%)	28(93.3%)
Yes	2(6.7%)	2(6.7%)

Table 1: Comparison of rescue analgesia- Group B patients needed rescue analgesics more often than Group A.

Complications	Group A (n=30)	Group B (n=30)	P value
Nausea/Vomiting	0	4(13.3%)	0.112
Hypotension	9(30.0%)	0	0.002**
Purities	0	5(16.7%)	0.052+
Resp Depression	0	0	-
Sedation	12(40.0%)	0	<0.001**
Bradycardia	0	0	-
Motor Blockade	5(16.7%)	1(3.3%)	0.195

Table 2: Comparison of Complications/Side effects between two groups of patients studied

AUTHORS:

1. Rachana N. D.
2. Saraswathi N.
3. Srinivas Babu A.
4. Anup N.

PARTICULARS OF CONTRIBUTORS:

1. Senior Resident, Department of Anaesthesia, The Bangalore Hospital.
2. Assistant Professor, Department of Anaesthesia, BMCRI-SSH.
3. HOD, Department of Anaesthesia, The Bangalore Hospital.
4. Consultant, Department of Anaesthesia, The Bangalore Hospital.

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Rachana N. D,
#124, Beml Layout,
Basaveswara Nagar,
Bangalore-560079.
Email: rachanakiran84@gmail.com

Date of Submission: 05/07/2014.
Date of Peer Review: 06/07/2014.
Date of Acceptance: 24/07/2014.
Date of Publishing: 31/07/2014.